



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The following occultations were observed, using a power of about 85:

S. D. -8° 5994.	Disappearance, $9^{\text{h}} 39^{\text{m}} 16^{\text{s}}.0$ P. S. T.
$11 \pm$ mag. star.	" $9^{\text{h}} 56^{\text{m}} 03^{\text{s}}.0$
$11 \pm$ mag. star.	" $9^{\text{h}} 57^{\text{m}} 08^{\text{s}}.0$
S. D. -8° 5996.	" $9^{\text{h}} 58^{\text{m}} 05^{\text{s}}.5$
A star of $11 \pm$ mag.	" $10^{\text{h}} 04^{\text{m}} 23^{\text{s}}.7$
S. D. -7° 5900.	" $10^{\text{h}} 09^{\text{m}} 37^{\text{s}}.9$
12 to 13 mag. star.	" $10^{\text{h}} 18^{\text{m}} 34^{\text{s}} \pm 1^{\text{s}}.$
S. D. -7° 5907.	" $10^{\text{h}} 50^{\text{m}} 21^{\text{s}} \pm 1^{\text{s}}.$

Soon after totality commenced, I turned the telescope to the place of SWIFT's comet, but, as it was near the horizon, failed to see it. Just before totality ended, I again turned to the comet's place, and this time found it without difficulty. There was not sufficient time to adjust the micrometer and make an observation.

Its light was about the same as during the latter part of August, but its centre seemed a little more condensed.

C. D. PERRINE.

LICK OBSERVATORY,
September 25, 1895.

TOTAL ECLIPSE OF THE MOON, SEPTEMBER 3, 1895.

The total eclipse of the Moon was observed with a three-inch telescope with alt-azimuth mounting, and the times of contact were noted by a mean time chronometer. The first contact was not observed, and the last contact was observed with the naked eye.

The occultation of several stars was noted, but only one star was bright enough to be well observed.

Fifteen minutes after totality began, all the more prominent features of the Moon's surface could be distinguished easily in the telescope, and even in the middle of the eclipse *Tycho* and a few other craters could be seen clearly. To the naked eye, the contrast between the light and dark areas was well marked throughout the eclipse, the color varying from orange-red to a deep copper-red.

The following edge of the shadow was heavier and more sharply defined than the preceding.

The star occulted was estimated at 9.5 magnitude, in position angle 95° with reference to the Moon's centre. The star was

BD.—8°, 5996 (8.7 mag.). The time of disappearance was 9^h 57^m 58^s.3 P. S. T.

Beginning of totality, 9^h 6^m 28^s P. S. T.

End of totality, 10^h 47^m 35^s

Last contact with shadow, 11^h 54^m 00^s

R. G. AITKEN.

LICK OBSERVATORY,

September 4, 1895.

LUNAR ECLIPSE, SEPTEMBER 3, 1895.

Beginning of totality, 9^h 6^m 35^s P. S. T.

End of totality, 10^h 47^m 20^s.

Observations made without telescopic aid.

W. W. CAMPBELL.

NOTE ON THE MELTING OF THE POLAR CAPS OF *MARS*.

I suppose no fact concerning *Mars* has been better established than the one that the polar caps continue to decrease in size after the summer solstice on the planet has passed. The statement has recently been made, both in an astronomical journal and in the secular press, that the continued diminution of the cap after solstice proves that the maximum temperature on the planet occurs several months after summer solstice, and, therefore, that *Mars* has a "heat-storing atmosphere."

I believe that astronomers have always considered that *Mars* has *some* atmosphere, that the more or less extensive atmosphere is necessarily "heat-storing," and that the maximum temperature necessarily, therefore, occurs some time after summer solstice. But does the continued melting of the caps after solstice prove it? I think not.

Suppose that a given area *A* of a cap melts off before solstice, leaving a remnant of area *B* covering the polar region. The area *B* receives the same amount of direct solar heat before solstice that it does after. If this heat had no effect upon the area *B* until after summer solstice, then the point would be well taken. But such is not the case. There is abundant positive proof that extensive melting of the area *B* does occur before solstice; and that it should continue to melt after solstice does not prove that the maximum temperature occurs then, since the same amount of direct solar heat is received after solstice as before.